

661.566: 66.097.3:669.054.8

• • " " , • • , • • , • • ,

The possibility of the utilization of platinum containing wastes of nitric acid production is studied. Quantitative content of the wastes containing platinum group metals from the surface of boil-utilizator and absorption column is shown. In this article content of wastes with platinum group metals and choose sedimentator for metal platinum obtaining are made.

[1]  
( ) [2],  
[3, 4].  
32 , , 2,4  
, 640 .  
:

,  
 -  
 ,  
 ,  
 ,  
 -  
 -  
 .  
 0,716 ,  
 - , , -  
 , .  
 ,  
 " " ,  
 , , -  
 -  
 .  
 -  
 0,716 . 2400 -  
 .  
 - , 1.  
 1

	, % .
	50,43
	1,37
	16,87
	1,1
	0,16
	0,34
	0,1
	17,75
	0,05
	0,25


	0,92
	0,16

,  
 ,  
 , 2.  
 2

	, %
	11,35
	1,05
	9,98
	13,43
	4,21
	0,25
	0,03
	27,28
	0,18
	0,22
	0,06

.1 .2, , , -  
 . -  
 , .  
 , , -  
 .  
 , ,  
 , .

∴ ,  
 , , , ..  
 [5]  
 , , , , ,  
 ( , , , )  
 , , -



800...1000 °C

~ 10 %

[6].

[7].

: 1. 1985. – 327 . 2. / 1999 – 368 . 3. . – . 1983. – 286 . 4. , 1974. - 17. - . 10. - . 1587 – 1589. 5. , 1972 – 613 . 6. // 2 " . – 1997. – . 1. – . 189-190. 7. . , 1965 – 215 .

16.04.07